



Centromere Protein-F-like Pattern in a Patient With Rheumatoid Arthritis

Kyuhwa Hur, M.D.¹, La-He Jearn, M.D.^{1,2}, and Think-You Kim , M.D.²

¹Department of Laboratory Medicine, Hanyang University Medical Center, Seoul, Korea; ²Department of Laboratory Medicine, College of Medicine, Hanyang University, Seoul, Korea

Dear Editor,

Autoantibodies to centromere protein-F (CENP-F), which is closely associated with cell proliferation, are known to be a specific marker for malignant tumors [1-3]. However, a few cases of autoantibodies to CENP-F have been reported in autoimmune diseases. Despite some studies indicating no association between autoimmune diseases and CENP-F [2, 3], the association still remains unclear. Particularly, there are no reports on autoantibodies to CENP-F or CENP-F-like pattern by antinuclear antibody (ANA) test in rheumatoid arthritis (RA). We observed the first case of a CENP-F-like pattern in RA without any underlying diseases.

A 37-year-old female visited Hanyang University Medical Center, Seoul, Korea, with 3-month history of pain in both knees and right ankle. The laboratory test results were as follows: erythrocyte sedimentation rate (ESR), 74 mm/hr; C-reactive protein (CRP), 4.4 mg/dL; rheumatoid factor (RF), 89 IU/mL; and anti-cyclic citrullinated peptide (anti-CCP), 176 U/mL. X-ray imaging of the hands and feet revealed soft tissue swelling, and whole-body bone scanning revealed abnormally increased bone uptake in the right foot, both hands, and both knees. The ANA test showed a CENP-F-like pattern and the titer of 1:640 (Fig. 1). All other findings were negative, with no underlying disease, family history, or drug history. She was diagnosed as having RA on the

basis of the 2010 American College of Rheumatology/European League Against Rheumatism RA classification criteria [4] with a total score of 7: two small and two large joints involved (score 2), high positive RF >42 IU/mL and anti-CCP >30 U/mL (score 3), abnormal ESR >20 mm/hr in females and CRP >0.8 mg/dL (score 1), and >6 weeks of symptoms (score 1).

CENP-F is a 367 kDa nuclear protein first reported in 1993 [5, 6]. It is distributed in the nuclear matrix at the early G₂ phase, forms a kinetochore at the late G₂ phase to promote activation of the centromere and cell division, and disappears after the completion of M phase. Perhaps, it promotes cell proliferation, an increase in the number of mitotic cells as the cell cycle gets faster, because it is specific to malignant tumors, breast cancer, lung cancer, ovarian cancer, cervical cancer, non-Hodgkin lymphoma, and esophageal squamous cell carcinoma [1-3, 6]. Particularly, the high expression level of CENP-F in primary breast cancer is considered a molecular background of the rapid proliferation and high aggressiveness of cancer cells [7].

The ANA test is the best method to detect fluorescence of a CENP-F-like pattern, which is expressed during the cell cycle (G₂-M phase). It appears similar to the nuclear speckled pattern at the early G₂ phase, multiple bright paired foci inside the nucleus and a smooth nuclear envelope similar to the anti-prolifer-

Received: March 13, 2018

Revision received: August 22, 2018

Accepted: October 16, 2018

Corresponding author: Think-You Kim, M.D.

 <https://orcid.org/0000-0002-4131-0107>

Departments of Laboratory Medicine and Diagnostic Immunology,
Hanyang University Medical Center, 222-1 Wangsimni-ro, Seongdong-gu,
Seoul 04763, Korea

Tel: +82-2-2290-8980, Fax: +82-2-2290-9193

E-mail: tykim@hanyang.ac.kr

© Korean Society for Laboratory Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1. The antinuclear antibody test of HEp-2 cells using the indirect immunofluorescence method. The image shows the centromere protein-F-like pattern. It comprises the nuclear speckled pattern, multiple bright paired foci inside the nucleus at the early G₂ phase; smooth nuclear envelope at the late G₂ phase; centromere pattern, multiple aligned small and faint dots at the prophase and metaphase; and midbody pattern, an intense dot located at the midzone at the late anaphase and telophase.

ating cell nuclear antigen (anti-PCNA) antibody pattern at the late G₂ phase. The centromere pattern, multiple aligned small dots, appears from prophase to metaphase; the midbody pattern, an intense dot located at the midzone, appears from late anaphase to telophase [2, 5, 6].

Several studies have reported autoantibodies to CENP-F; however, none have explained their correlation with RA. In previous studies, patients with malignant tumors and autoantibodies to CENP-F did not exhibit any relationship with RA [2, 3]. Moreover, six patients with autoantibodies to CENP-F did not have RA (undefined connective tissue disease, 2; primary antiphospholipid syndrome, 1; colorectal carcinoma, 1; hepatitis C virus, 1; and fever of unknown origin, 1) [8]. Another six patients with both RA and malignant tumors were reported without any correlation with CENP-F [9].

As our patient's symptoms and laboratory tests did not suggest malignant tumors, the likelihood of malignancy was low, and the CENP-F-like pattern was related to RA rather than malignancy. However, no case having a CENP-F-like pattern in RA without any underlying diseases has been reported thus far, and the correlation between CENP-F and RA is not clear. Temporary acute inflammatory reaction to RA might account for an increase in autoantibodies to CENP-F. Therefore, periodic ANA tests to continuously detect a CENP-F-like pattern are imperative.

Several patterns of the ANA test, including speckled, homogeneous, and centromere patterns, are commonly reported in RA [10]. The CENP-F-like pattern might be misinterpreted as the speckled, centromere, or anti-PCNA antibody pattern. We think that CENP-F-like patterns in RA have been incorrectly reported and dismissed. To date, the importance of autoantibodies to CENP-F in RA remains understated. Therefore, if more cases of CENP-F-like patterns in RA are reported, further studies on its clinical relevance in RA are warranted.

Authors' Disclosures of Potential Conflicts of Interest

No potential conflicts of interest relevant to this article were reported.

REFERENCES

1. Fritzier MJ, Rattner JB, Luft LM, Edworthy SM, Casiano CA, Peebles C, et al. Historical perspectives on the discovery and elucidation of autoantibodies to centromere proteins (CENP) and the emerging importance of antibodies to CENP-F. *Autoimmun Rev* 2011;10:194-200.
2. Casiano CA, Humbel RL, Peebles C, Covini G, Tan EM. Autoimmunity to the cell cycle-dependent centromere protein p330d/CENP-F in disorders associated with cell proliferation. *J Autoimmun* 1995;8:575-86.
3. Rattner JB, Rees J, Whitehead CM, Casiano CA, Tan EM, Humbel RL, et al. High frequency of neoplasia in patients with autoantibodies to centromere protein CENP-F. *Clin Invest Med* 1997;20:308-19.
4. Aletaha D, Neogi T, Silman AJ, Funovits J, Felson DT, Bingham CO 3rd, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum* 2010;62:2569-81.
5. Liao H, Winkfein RJ, Mack G, Rattner JB, Yen TJ. CENP-F is a protein of the nuclear matrix that assembles onto kinetochores at late G₂ and is rapidly degraded after mitosis. *J Cell Biol* 1995;130:507-18.
6. Casiano CA, Landberg G, Ochs RL, Tan EM. Autoantibodies to a novel cell cycle-regulated protein that accumulates in the nuclear matrix during S phase and is localized in the kinetochores and spindle midzone during mitosis. *J Cell Sci* 1993;106:1045-56.
7. Ueda S, Kondoh N, Tsuda H, Yamamoto S, Asakawa H, Fukatsu K, et al. Expression of centromere protein F (CENP-F) associated with higher FDG uptake on PET/CT, detected by cDNA microarray, predicts high-risk patients with primary breast cancer. *BMC Cancer* 2008;8:384.
8. Bonaci-Nikolic B, Andrejevic S, Bukilica M, Urosevic I, Nikolic M. Autoantibodies to mitotic apparatus: association with other autoantibodies and their clinical significance. *J Clin Immunol* 2006;26:438-46.
9. Wen J, Ouyang H, Yang R, Bo L, Zhang Y, Tang M, et al. Malignancy dominated with rheumatic manifestations: a retrospective single-center analysis. *Sci Rep* 2018;8:1786.
10. Garcia-de la Torre I and Miranda-Mendez L. Studies of antinuclear antibodies in rheumatoid arthritis. *J Rheumatol* 1982;9:603-6.