



# Rejuvenating Aged Hematopoietic Stem Cells Through Improvement of Mitochondrial Function

James Moon, M.S.<sup>1,2</sup>, Hye Ran Kim , Ph.D.<sup>3</sup>, and Myung Geun Shin , M.D.<sup>1,4,5</sup>

Department of Laboratory Medicine<sup>1</sup>, Chonnam National University Medical School and Chonnam National University Hwasun Hospital, Jeollanam-do, Korea; Department of Biomedical Engineering<sup>2</sup>, University of California, CA, USA; College of Korean Medicine<sup>3</sup>, Dongshin University, Naju, Korea; Brain Korea 21 Plus Project<sup>4</sup>, Chonnam National University Medical School, Gwangju, Korea; Environmental Health Center for Childhood Leukemia and Cancer<sup>5</sup>, Chonnam National University Medical School and Chonnam National University Hwasun Hospital, Jeollanam-do, Korea

Mitochondria are the powerhouses of the cell as well as the primary site of hematopoiesis, which also occurs in the cytoplasm. Hematopoietic stem cells (HSCs) are characterized by a very high turnover rate, and are thus considered to be relatively free from the age-related insults generated by mitochondria. However, HSCs are also subject to these age-related insults, including the incidence of myeloid proliferative diseases, marrow failure, hematopoietic neoplasms, and deterioration of the adaptive human immune system. Recently, NAD<sup>+</sup> dietary supplements, known as niacin or vitamin B<sub>3</sub>, including tryptophan, nicotinic acid, nicotinamide, and the newly identified NAD<sup>+</sup> precursor nicotinamide riboside, have been shown to play a role in restoring adult stem cell function through the amelioration of mitochondrial dysfunction. This insight motivated a study that focused on reversing aging-related cellular dysfunction in adult mouse muscle stem cells by supplementing their diet with nicotinamide riboside. The remedial effect of nicotinamide riboside enhanced mitochondrial function in these muscle stem cells in a SIRT1-dependent manner, affecting cellular respiration, membrane potential, and production of ATP. Accordingly, numerous studies have demonstrated that sirtuins, under nuclear/mitochondrial control, have age-specific effects in determining HSC phenotypes. Based on the evidence accumulated thus far, we propose a clinical intervention for the restoration of aged HSC function by improving mitochondrial function through NAD<sup>+</sup> precursor supplementation.

**Key Words:** Hematopoietic stem cells, Aging, Mitochondrial dysfunction, Restoration, NAD<sup>+</sup> precursors

**Received:** April 30, 2017  
**Revision received:** July 24, 2017  
**Accepted:** May 4, 2018

**Corresponding author:** Hye Ran Kim  
 <https://orcid.org/0000-0002-0658-3345>  
College of Korean Medicine, Dongshin University, 185 Gunjae-ro, Naju, Jeollanam-do 58245, Korea  
Tel: +82-61-330-3514  
Fax: +82-61-330-3519  
E-mail: 98lani@gmail.com

**Co-corresponding author:** Myung Geun Shin  
 <https://orcid.org/0000-0002-0372-9185>  
Department of Laboratory Medicine, Chonnam National University Hwasun Hospital, 322 Seoyang-ro, Hwasun-eup, Hwasun-gun, Jeollanam-do 58128, Korea  
Tel: +82-61-379-7950  
Fax: +82-61-379-7984  
E-mail: mgshin@jnu.ac.kr

**© Korean Society for Laboratory Medicine**

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Mitochondria are generally characterized as the powerhouse of the cell, since this is the site where energy is produced from ATP. In addition to energy production, mitochondria play a key role in several important cellular processes, including growth, signaling, differentiation, reactive oxygen species (ROS) production, apoptosis, and cell cycle control. Interestingly, unlike other cellular organelles, mitochondria have their own DNA, mito-

chondrial DNA (mtDNA), and several studies have indicated an association between the accumulation of mtDNA mutations and mammalian aging [1-3].

Historically, mitochondria have not been considered important in restoring the functions of aged hematopoietic stem cells (HSCs); however, emerging studies on rejuvenating HSCs suggest an association between sirtuins (SIRT) and mitochondrial activities [4, 5]. In addition, a study on the deregulation of the mitochondrial stress-mediated metabolic system demonstrated that

SIRT7 strongly influences the regenerative capacity of HSCs [6]. Although the functions of musculoskeletal stem cells (MuSCs) and HSCs are distinct, alteration of the SIRT1-associated nuclear/mitochondrial axis appears to be a common hallmark of aging in both cell types [7, 8].

Recent research suggests the possibility of restoring the mitochondrial functions of aged stem cells, including MuSCs, nerve tissue stem cells (NSCs), and melanocyte stem cells (McSCs), by NAD<sup>+</sup> supplementation without genetic manipulation [8, 9]. The remedial effect of the NAD<sup>+</sup> precursor nicotinamide riboside (NR) enhances mitochondrial functions in stem cells, including respiration, membrane potential, ATP production, and the mitochondrial unfolded protein response (UPR); however, these effects are not observed in stem cells with a SIRT1 deficit. Moreover, NR was found to suppress the process of senescence in adult NSCs and McSCs [8].


These findings have reinforced the notion that NAD<sup>+</sup> precursors can function as a pharmacological tool to enhance SIRT activities. This, in turn, paves the way for clinical translation of NAD<sup>+</sup> precursor treatment through further investigations of hematopoietic tissues. We review evidence relating mitochondrial dysfunction to HSC aging, and propose a strategy for mitochondrial-targeted recovery as a potentially safe, effective, and non-invasive method for the control or prevention of aging-related hematopoietic diseases.

## ROLE OF THE MITOCHONDRIA IN HUMAN HEMATOPOIESIS

Mitochondria are central to the heme biosynthetic pathway, part of which occurs in the cytoplasm, eventually returning to the mitochondrion. Enzyme defects in the heme biosynthetic pathway cause sideroblastic anemia, leading to a deficiency of heme precursors and mitochondria that cannot fully utilize iron. In erythroid precursors, most of the iron initially gains access to the cell through transferring receptors, subsequently entering the mitochondria where it combines with protoporphyrin IX to produce heme. The heme produced leaves the mitochondria to merge with globin chains and synthesize cytoplasmic ribosomes [10]. Thus, when protoporphyrin synthesis is highly impaired, the imported iron instead accumulates in the mitochondria due to reactant deficiency (Fig. 1).

## CONTRIBUTION OF MITOCHONDRIAL DYSFUNCTION AND GENOME ALTERATIONS TO HSC AGING

HSCs are generally dormant but have the potential to become highly active to restore blood on demand. Sustenance of HSC dormancy requires supply of low metabolic activity by glycolytic metabolites [11, 12]. Thus, unlike MuSCs, NSCs, and McSCs,



**Fig. 1.** The heme biosynthetic pathway in the mitochondria and cytoplasm. Heme synthesis starts in the mitochondria and then returns to the mitochondria through several cytoplasmic steps. Abbreviation: ALA, aminolevulinic acid.


which highly depend on mitochondrial ATP generation, the role of mitochondria in HSC homeostasis has traditionally not been emphasized.

During ATP production through oxidative phosphorylation, ROS are produced as the by-product of mitochondrial respiration [13]. Owing to their low metabolic activity, dormant HSCs exhibit very low levels of ROS that are closely associated with cellular metabolic activity [12, 14]. Accordingly, this unbalanced accumulation of ROS mediates HSC dysfunction [15-17], and recent evidence implies that mitochondria play crucial roles in the maintenance of HSC quiescence and their capacity to switch from dormancy to a metabolically active state [18-21].

From a clinical perspective, HSCs are characterized by a very high turnover rate; however, they are not exempt from age-related insults. Aged HSCs are associated with increased incidence of myeloid proliferative diseases, such as marrow failure and hematopoietic neoplasms, and deterioration of the adaptive human immune system [22]. Because HSCs play a crucial role in maintaining the circulation, functional deterioration of HSCs may be highly responsible for age-related damage. Although HSCs maintain hematopoiesis for multiple processes, they are

subject to drastic phenotypic and functional changes during aging, as confirmed by serial transplantation studies in mice [22]. The most notable of these changes is failure of the adaptive immune system, resulting in the weakened lymphoid function that is common in the elderly. In addition, aging often leads to the overproduction of myeloid cells, which fosters a pro-inflammatory hematopoietic environment [22, 23].

Aging leads to several clinical conditions related to the hematopoietic system, including decreased functionality of the adaptive immune system, elevated incidences of certain autoimmune diseases, age-associated anemia, and hematological malignancies [22]. Similar to other human tissues, the aged hematopoietic system experiences a decline in regenerative capacity for normal homeostasis after stress or damage [22]. Several pathways are involved in the aging mechanisms of the hematopoietic system, and both intrinsic and extrinsic factors are related to the aging environment (Fig. 2). However, recent studies suggest that age-dependent cellular and mitochondrial damage within the most primitive HSCs may play a crucial role in hematopoietic deterioration during aging [24, 25].



**Fig. 2.** Proposed molecular mechanisms of hematopoietic stem cell (HSC) aging and related disease phenotypes. HSC aging is caused by the alteration of nuclear and mitochondrial genomes, as well as transcriptome and epigenetic changes, resulting in the occurrence of aging-related hematopoietic diseases.

Abbreviations: mtDNA, mitochondrial DNA; ROS, reactive oxygen species.

## REJUVENATING AGED HSCS THROUGH RESTORING MITOCHONDRIAL DYSFUNCTION

One possible explanation for mammalian mitochondrial dysfunction during aging is the accumulation of altered nuclear and mitochondrial genetic materials, although environmental factors are clearly at play as well.

NAD was discovered over a century ago, and its role as a redox agent in metabolism has subsequently been established [26]. More recently, the oxidized form, NAD<sup>+</sup>, was revealed to be a key factor in mitochondrial function, and NAD<sup>+</sup> supplementation has been shown to restore the normal phenotype [26]. Recent studies have clearly shown the role of NAD<sup>+</sup> dietary supplements, commonly referred to as niacin or vitamin B<sub>3</sub>, in mitochondrial function, including tryptophan (Trp), nicotinic acid (Na), nicotinamide (Nam), and the newly identified NAD<sup>+</sup> precursor NR [27-30].

More importantly, aging is related to SIRT deficiency and decreased mitochondrial function, and the NAD<sup>+</sup>/SIRT pathway is a pivotal factor in sustaining health and forestalling age-related diseases [31]. Indeed, SIRT influence a broad range of cellular activities such as aging, transcription, apoptosis, and inflammation [32]. One study, using engineered mice expressing excess SIRT1, demonstrated that the level of cellular NAD<sup>+</sup> gradually drops during normal aging [33]. Furthermore, NR was shown to enhance oxidative metabolism by increasing the NAD<sup>+</sup> level and activating SIRT1 and SIRT3, suggesting the potential of NR as a pharmacological supplement to recover the metabolic and age-related disorders characterized by mitochondrial dysfunction [34].


Many studies have provided evidence indicating that SIRT of

the nucleus/mitochondrial control axis have age-specific effects in mediating the phenotypes of HSCs. Differentiation of SIRT1-deleted HSCs exhibited the typical characteristics of aged HSCs, such as a decline in the lymphoid compartment, anemia, and altered expression of related genes, indicating an essential role of SIRT1 in HSC homeostasis [7]. SIRT3 regulates the global acetylation of mitochondrial proteins and a stress response. Brown *et al* [4] showed that SIRT3 expression was suppressed with aging, and upregulation of SIRT3 enhanced the regenerative capacity of aged HSCs. When SIRT7, which controls the deacetylation promoter, was inactivated in HSCs, quiescence was reduced and the mitochondrial protein folding stress (PFS<sup>mt</sup>) occurred, ultimately resulting in compromised regenerative capacity. By contrast, SIRT7 up-regulation enhanced the regenerative capacity of aged HSCs [6]. Another recent study also showed that SIRT6-deficient HSCs exhibited impaired self-renewal capacity [27].

## POSSIBLE APPROACH FOR CORRECTING AGING-RELATED HSC DYSFUNCTION

Based on the studies described above, we propose a mitochondria-targeted strategy for controlling the HSC aging mechanism and associated regulatory factors toward restoration of aged HSC function through improving mitochondrial function (Fig. 3).


The aging process of HSCs is potentially linked with poly ADP-ribose polymerase (PARP) activation, NAD<sup>+</sup> deficiency, SIRT inactivation, mitochondrial dysfunction, and cell and tissue damage, reflecting a generalized aging syndrome, which may be corrected by supplementation with NAD<sup>+</sup> precursors (Fig. 4). In particular, recent studies have indicated that NR supplementa-




**Fig. 3.** Schematic representation of the research objectives and hypothesis: restoration of aged hematopoietic stem cell (HSC) function through improving mitochondrial function.

tion could rejuvenate aged stem cells such as MuSCs, NSCs, and McSCs, enhance lifespan, and improve muscle function in

muscular dystrophy models [8, 9]. Moreover, the association of SIRT1 with the aging-like phenotypes of both MuSCs and HSCs



**Fig. 4.** Aging syndrome of a hematopoietic stem cell (HSC) and proposed restoration strategy. In old animals, aging may cause an NAD<sup>+</sup> deficit by activating poly (ADP-ribose) polymerases (PARPs). The NAD<sup>+</sup> shortage then decreases the activity of sirtuins, which are anti-aging proteins, thereby promoting the aging cycle. Nicotinamide riboside (NR) replenishes NAD<sup>+</sup> and corrects the cycle to enhance stem cell maintenance, tissue function, and the unfolded protein response (UPR).



**Fig. 5.** Possible molecular mechanisms and role of NAD<sup>+</sup> precursors in restoring hematopoietic stem cell (HSC) function. Aged HSC function might be recovered through increasing mitochondrial function using NAD<sup>+</sup> precursors. Abbreviations: AMPK, 5' adenosine monophosphate-activated protein kinase; NAMPT, nicotinamide phosphoribosyltransferase; FOXO, forkhead box protein; SIRT1, sirtuin 1; Sir2, silent information regulator 2; NR, nicotinamide riboside; cADP, cyclic ADP; Nam, nicotinamide; Trp, tryptophan; Na, nicotinic acid; Na ADP, nicotinic ADP.

suggests a plausible pharmacological approach by targeting the interplay of SIRT6 and the nuclear/mitochondrial control axis in HSCs [4, 5, 7, 8].

An important consideration is to examine the feasibility of activating SIRT3, SIRT6, and SIRT7, which have specific rejuvenation effects on HSCs [4-6]. For example, experiments with SIRT1-deficient mice provided NR supplementation in tablet form showed improvement of muscular dystrophy through enhancing mitochondrial function [8, 9]. This implies that NR would easily access human hematopoietic tissues to exert the expected rejuvenation effect of aged HSCs.

Moreover, NR supplementation in mammalian cells increases NAD<sup>+</sup> levels and activates SIRT1 and SIRT3, leading to amelioration of the metabolic and age-related degeneration characterized by mitochondrial dysfunction [34]. Therefore, more detailed studies on the effects of NR, the NAD<sup>+</sup> precursor, on SIRT6 to improve the mitochondrial activities of HSCs may lead to the development of optimal strategies for reversing age-dependent degeneration with an accessible approach.

Thus, NAD<sup>+</sup> precursors could improve mitochondrial function to recover aged HSC function through the hypothetical pathway shown in Fig. 5 [35, 36].

## CONCLUSION

Although further research on the effects of NAD<sup>+</sup> replenishment in human health maintenance is required, the studies we reviewed clearly demonstrate that the level of NAD<sup>+</sup> drops during the aging process [37]. In addition, the relevant pathways of NAD<sup>+</sup> synthesis, SIRT6, and PARPs suggest that NAD<sup>+</sup> replenishment may be beneficial in rejuvenating aged HSCs.

The ability to rejuvenate stem cells without relying on genetic manipulation is the safest way to achieve optimal clinical outcomes. Regulation of aging through enhancing mitochondrial function is a potentially effective, low-cost, and stable treatment method.

## Authors' Disclosure of Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

## Acknowledgements

This study was supported by a National Research Foundation of Korea (NRF) grant (No. 2016-R1C1B106130) funded by the

Korean Ministry of Science, ICT and Future Planning; the Leading Foreign Research Institute Recruitment Program (No. 2011-0030034) through the NRF funded by the Ministry of Education, Science and Technology (MEST); and the Environmental Health Center funded by the Ministry of Environment, Korea.

## REFERENCES

1. Kujoth G, Hiona A, Pugh TD, Someya S, Panzer K, Wohlgemuth SE, et al. Mitochondrial DNA mutations, oxidative stress, and apoptosis in mammalian aging. *Science* 2005;309:481-4.
2. Trifunovic A, Wredenberg A, Falkenberg M, Spelbrink JN, Rovio AT, Bruder CE, et al. Premature ageing in mice expressing defective mitochondrial DNA polymerase. *Nature* 2004;429:417-23.
3. Bratic A and Larsson NG. The role of mitochondria in aging. *J Clin Invest* 2013;123:951-7.
4. Brown K, Xie S, Qiu X, Mohrin M, Shin J, Liu Y, et al. SIRT3 reverses aging-associated degeneration. *Cell Rep* 2013;3:319-27.
5. Wang H, Diao D, Shi Z, Zhu X, Gao Y, Gao S, et al. SIRT6 controls hematopoietic stem cell homeostasis through epigenetic regulation of Wnt signaling. *Cell Stem Cell* 2016;18:495-507.
6. Mohrin M, Shin J, Liu Y, Brown K, Luo H, Xi Y, et al. A mitochondrial UPR-mediated metabolic checkpoint regulates hematopoietic stem cell aging. *Science* 2015;347:1374-7.
7. Rimmelé P, Bigarella CL, Liang R, Izac B, Diequez-Gonzalez R, Barbet G, et al. Aging-like phenotype and defective lineage specification in SIRT1-deleted hematopoietic stem and progenitor cells. *Stem Cell Rep* 2014;3:44-59.
8. Zhang H, Ryu D, Wu Y, Gariani K, Wang X, Luan P, et al. NAD<sup>+</sup> repletion improves mitochondrial and stem cell function and enhances life span in mice. *Science* 2016;352:1436-43.
9. Ryu D, Zhang H, Ropelle ER, Sorrentino V, Mázala DA, Mouchiroud L, et al. NAD<sup>+</sup> repletion improves muscle function in muscular dystrophy and counters global PARylation. *Science Transl Med* 2016;8:361ra139.
10. Gattermann N. From sideroblastic anemia to the role of mitochondrial DNA mutations in myelodysplastic syndromes. *Leuk Res* 2000;24:141-51.
11. Unwin RD, Smith DL, Blinco D, Wilson CL, Miller CJ, Evans CA, et al. Quantitative proteomics reveals posttranslational control as a regulatory factor in primary hematopoietic stem cells. *Blood* 2006;107:4687-94.
12. Simsek T, Kocabas F, Zheng J, DeBerardinis RJ, Mahmoud AI, Olsen EN, et al. The distinct metabolic profile of hematopoietic stem cells reflects their location in a hypoxic niche. *Cell Stem Cell* 2010;7:380-90.
13. Finkel T. Signal transduction by mitochondrial oxidants. *J Biol Chem* 2012;287:4434-40.
14. Jang YY and Sharkis SJ. A low level of reactive oxygen species selects for primitive hematopoietic stem cells that may reside in the low-oxygenic niche. *Blood* 2007;110:3056-63.
15. Ito K, Hirao A, Arai F, Matsuoka S, Takubo K, Hamaguchi I, et al. Regulation of oxidative stress by ATM is required for self-renewal of haematopoietic stem cells. *Nature* 2004;431:997-1002.
16. Kocabas F, Zheng J, Thet S, Copeland NG, Jenkins NA, DeBerardinis RJ, et al. Meis1 regulates the metabolic phenotype and oxidant defense of hematopoietic stem cells. *Blood* 2012;120:4963-72.
17. Zheng J, Lu Z, Kocabas F, Böttcher RT, Costell M, Kang X, et al. Profilin 1 is essential for retention and metabolism of mouse hematopoietic stem cells in bone marrow. *Blood* 2014;123:992-1001.

18. Chen C, Liu Y, Liu R, Ikenoue T, Guan KL, Liu Y, et al. TSC–mTOR maintains quiescence and function of hematopoietic stem cells by repressing mitochondrial biogenesis and reactive oxygen species. *J Exp Med* 2008;205:2397-408.
19. Nakada D, Saunders TL, Morrison SJ. Lkb1 regulates cell cycle and energy metabolism in haematopoietic stem cells. *Nature* 2010;468:653-8.
20. Norddahl GL, Pronk CJ, Wahlestedt M, Sten G, Nygren JM, Ugale A, et al. Accumulating mitochondrial DNA mutations drive premature hematopoietic aging phenotypes distinct from physiological stem cell aging. *Cell Stem Cell* 2011;8:499-510.
21. Takubo K, Nagamatsu G, Kobayashi CI, Nakamura-Ishizu A, Kobayashi H, Ikeda E, et al. Regulation of glycolysis by Pdk functions as a metabolic checkpoint for cell cycle quiescence in hematopoietic stem cells. *Cell Stem Cell* 2013;12:49-61.
22. Ergen AV and Goodell MA. Mechanisms of hematopoietic stem cell aging. *Exp Gerontol* 2010;45:286-90.
23. Park CB and Larsson NG. Mitochondrial DNA mutations in disease and aging. *J Cell Biol* 2011;193:809-18.
24. Shin MG, Kajigaya S, McCoy JP Jr, Levin BC, Young NS. Marked mitochondrial DNA sequence heterogeneity in single CD34+ cell clones from normal adult bone marrow. *Blood* 2004;103:553-61.
25. Kim HR, Won SJ, Fabian C, Kang MG, Szardenings M, Shin MG. Mitochondrial DNA aberrations and pathophysiological implications in hematopoietic diseases, chronic inflammatory diseases, and cancers. *Ann Lab Med* 2015;35:1-14.
26. Gomes AP, Price NL, Ling AJ, Moslehi JJ, Montgomery MK, Rajman L, et al. Declining NAD+ induces a pseudohypoxic state disrupting nuclear-mitochondrial communication during aging. *Cell* 2013;155:1624-38.
27. Lin J, Pan Y, Wang J. NAD+ and its precursors in human longevity. *Quant Biol* 2015;3:193-8.
28. Bieganski P and Brenner C. Discoveries of nicotinamide riboside as a nutrient and conserved NRK genes establish a Preiss-Handler independent route to NAD+ in fungi and humans. *Cell* 2004;117:495-502.
29. Belenky P, Racette FG, Bogan KL, McClure JM, Smith JS, Brenner C. Nicotinamide riboside promotes Sir2 silencing and extends lifespan via Nrk and Urh1/Pnp1/Meu1 pathways to NAD+. *Cell* 2007;129:473-84.
30. Belenky P, Bogan KL, Brenner C. NAD+ metabolism in health and disease. *Trends Biochem Sci* 2007;32:12-9.
31. van de Ven RAH, Santos D, Haigis MC. Mitochondrial sirtuins and molecular mechanisms of aging. *Trends Mol Med* 2017;23:320-331.
32. Preyat N and Leo O. Sirtuin deacylases: a molecular link between metabolism and immunity. *J Leuk Biol* 2013;93:669-80.
33. Ramsey KM, Milis KF, Satoh A, Imai S. Age-associated loss of Sirt1-mediated enhancement of glucose-stimulated insulin secretion in beta cell-specific Sirt1-overexpressing (BESTO) mice. *Aging Cell* 2008;7:78-88.
34. Cantó C, Houtkooper RH, Pirinen E, Youn DY, Oosterveer MH, Cen Y, et al. The NAD(+) precursor nicotinamide riboside enhances oxidative metabolism and protects against high-fat diet-induced obesity. *Cell Metab* 2012;15:838-47.
35. Cantó C and Auwerx J. Caloric restriction, SIRT1 and longevity. *Trends Endocrinol Metab* 2009;20:325-31.
36. Bogan KL and Brenner C. Nicotinic acid, nicotinamide, and nicotinamide riboside: a molecular evaluation of NAD+ precursor vitamins in human nutrition. *Annu Rev Nutr* 2008;28:115-30.
37. Massudi H, Grant R, Braidy N, Guest J, Farnsworth B, Guillemin GJ. Age-associated changes in oxidative stress and NAD+ metabolism in human tissue. *PLoS One* 2012;7:e42357.